

COMP2610/COMP6261 - Information Theory

Lecture 7: Relative Entropy and Mutual Information

Mark Reid and **Aditya Menon**

Research School of Computer Science
The Australian National University



August 12th, 2014

- Information content and entropy: definition and computation
- Entropy and average code length
- Entropy and minimum expected number of binary questions
- Joint and conditional entropies, chain rule

Let X be a random variable with outcomes in \mathcal{X}

Let $p(x)$ denote the probability of the outcome $x \in \mathcal{X}$

The (Shannon) information content of outcome x is

$$h(x) = \log_2 \frac{1}{p(x)}$$

As $p(x) \rightarrow 0$, $h(x) \rightarrow +\infty$ (rare outcomes are more informative)

Entropy: Review

The entropy is the average information content of all outcomes:

$$H(X) = \sum_x p(x) \log_2 \frac{1}{p(x)}$$

Entropy is minimised if \mathbf{p} is peaked, and maximized if \mathbf{p} is uniform:

$$0 \leq H(X) \leq \log|\mathcal{X}|$$

Entropy is related to minimal number of bits needed to describe a random variable

This time

- The decomposability property of entropy
- Relative entropy and divergences
- Mutual information

- 1 Decomposability of Entropy
- 2 Relative Entropy / KL Divergence
- 3 Mutual Information
 - Definition
 - Joint and Conditional Mutual Information
- 4 Wrapping up

Decomposability of Entropy

Example 1 (Mackay, 2003)

Let $X \in \{0, 1, 2\}$ be a r.v. created by the following process:

- 1 Flip a fair coin to determine whether $X = 0$

Decomposability of Entropy

Example 1 (Mackay, 2003)

Let $X \in \{0, 1, 2\}$ be a r.v. created by the following process:

- 1 Flip a fair coin to determine whether $X = 0$
- 2 If $X \neq 0$ flip another fair coin to determine whether $X = 1$ or $X = 2$

Decomposability of Entropy

Example 1 (Mackay, 2003)

Let $X \in \{0, 1, 2\}$ be a r.v. created by the following process:

- 1 Flip a fair coin to determine whether $X = 0$
- 2 If $X \neq 0$ flip another fair coin to determine whether $X = 1$ or $X = 2$

The probability distribution of X is given by:

$$p(X = 0) =$$

$$p(X = 1) =$$

$$p(X = 2) =$$

Decomposability of Entropy

Example 1 (Mackay, 2003)

Let $X \in \{0, 1, 2\}$ be a r.v. created by the following process:

- 1 Flip a fair coin to determine whether $X = 0$
- 2 If $X \neq 0$ flip another fair coin to determine whether $X = 1$ or $X = 2$

The probability distribution of X is given by:

$$p(X = 0) = \frac{1}{2}$$

$$p(X = 1) =$$

$$p(X = 2) =$$

Decomposability of Entropy

Example 1 (Mackay, 2003)

Let $X \in \{0, 1, 2\}$ be a r.v. created by the following process:

- 1 Flip a fair coin to determine whether $X = 0$
- 2 If $X \neq 0$ flip another fair coin to determine whether $X = 1$ or $X = 2$

The probability distribution of X is given by:

$$p(X = 0) = \frac{1}{2}$$

$$p(X = 1) = \frac{1}{4}$$

$$p(X = 2) =$$

Decomposability of Entropy

Example 1 (Mackay, 2003)

Let $X \in \{0, 1, 2\}$ be a r.v. created by the following process:

- 1 Flip a fair coin to determine whether $X = 0$
- 2 If $X \neq 0$ flip another fair coin to determine whether $X = 1$ or $X = 2$

The probability distribution of X is given by:

$$p(X = 0) = \frac{1}{2}$$

$$p(X = 1) = \frac{1}{4}$$

$$p(X = 2) = \frac{1}{4}$$

Decomposability of Entropy

Example 1 (Mackay, 2003) — Cont'd

By definition,

$$H(X) = \frac{1}{2} \log 2 + \frac{1}{4} \log 4 + \frac{1}{4} \log 4 = 1.5 \text{ bits.}$$

But imagine learning the value of X *gradually*:

Decomposability of Entropy

Example 1 (Mackay, 2003) — Cont'd

By definition,

$$H(X) = \frac{1}{2} \log 2 + \frac{1}{4} \log 4 + \frac{1}{4} \log 4 = 1.5 \text{ bits.}$$

But imagine learning the value of X *gradually*:

- 1 First we learn whether $X = 0$:
 - ▶ Binary variable with $\mathbf{p}^{(1)} = (\frac{1}{2}, \frac{1}{2})$
 - ▶ Hence $H(1/2, 1/2) = \log_2 2 = 1$ bit.

Decomposability of Entropy

Example 1 (Mackay, 2003) — Cont'd

By definition,

$$H(X) = \frac{1}{2} \log 2 + \frac{1}{4} \log 4 + \frac{1}{4} \log 4 = 1.5 \text{ bits.}$$

But imagine learning the value of X *gradually*:

- 1 First we learn whether $X = 0$:
 - ▶ Binary variable with $\mathbf{p}^{(1)} = (\frac{1}{2}, \frac{1}{2})$
 - ▶ Hence $H(1/2, 1/2) = \log_2 2 = 1$ bit.
- 2 If $X \neq 0$ we learn the the value of the second coin flip:
 - ▶ Also binary variable with $\mathbf{p}^{(2)} = (\frac{1}{2}, \frac{1}{2})$
 - ▶ Therefore $H(1/2, 1/2) = 1$ bit.

Decomposability of Entropy

Example 1 (Mackay, 2003) — Cont'd

By definition,

$$H(X) = \frac{1}{2} \log 2 + \frac{1}{4} \log 4 + \frac{1}{4} \log 4 = 1.5 \text{ bits.}$$

But imagine learning the value of X *gradually*:

- 1 First we learn whether $X = 0$:
 - ▶ Binary variable with $\mathbf{p}^{(1)} = (\frac{1}{2}, \frac{1}{2})$
 - ▶ Hence $H(1/2, 1/2) = \log_2 2 = 1$ bit.
- 2 If $X \neq 0$ we learn the the value of the second coin flip:
 - ▶ Also binary variable with $\mathbf{p}^{(2)} = (\frac{1}{2}, \frac{1}{2})$
 - ▶ Therefore $H(1/2, 1/2) = 1$ bit.

However, the second revelation only happens half of the time:

$$H(X) = H(1/2, 1/2) + \frac{1}{2}H(1/2, 1/2) = 1.5 \text{ bits.}$$

Decomposability of Entropy

Generalization

For a r.v. with probability distribution $\mathbf{p} = (p_1, \dots, p_{|\mathcal{X}|})$:

$$H(\mathbf{p}) = H(p_1, 1 - p_1) + (1 - p_1)H\left(\frac{p_2}{1 - p_1}, \dots, \frac{p_{|\mathcal{X}|}}{1 - p_1}\right)$$

Decomposability of Entropy

Generalization

For a r.v. with probability distribution $\mathbf{p} = (p_1, \dots, p_{|\mathcal{X}|})$:

$$H(\mathbf{p}) = H(p_1, 1 - p_1) + (1 - p_1)H\left(\frac{p_2}{1 - p_1}, \dots, \frac{p_{|\mathcal{X}|}}{1 - p_1}\right)$$

$H(p_1, 1 - p_1)$ = entropy for a random variable corresponding to “Is $X = x_0$?”

$H\left(\frac{p_2}{1 - p_1}, \dots, \frac{p_{|\mathcal{X}|}}{1 - p_1}\right)$ = entropy for a random variable corresponding to outcomes when $X \neq x_0$

$(1 - p_1)$ = probability of $X \neq x_0$

Decomposability of Entropy

Generalization

In general, we have that for any m :

$$\begin{aligned} H(\mathbf{p}) &= H\left(\sum_{i=1}^m p_i, \sum_{i=m+1}^{|\mathcal{X}|} p_i\right) \\ &+ \left(\sum_{i=1}^m p_i\right) H\left(\frac{p_1}{\sum_{i=1}^m p_i}, \dots, \frac{p_m}{\sum_{i=1}^m p_i}\right) \\ &+ \left(\sum_{i=m+1}^{|\mathcal{X}|} p_i\right) H\left(\frac{p_{m+1}}{\sum_{i=m+1}^{|\mathcal{X}|} p_i}, \dots, \frac{p_{|\mathcal{X}|}}{\sum_{i=m+1}^{|\mathcal{X}|} p_i}\right) \end{aligned}$$

Apply this formula with $m = 1$, $|\mathcal{X}| = 3$, $\mathbf{p} = (p_1, p_2, p_3) = (1/2, 1/4, 1/4)$

- 1 Decomposability of Entropy
- 2 Relative Entropy / KL Divergence
- 3 Mutual Information
 - Definition
 - Joint and Conditional Mutual Information
- 4 Wrapping up

Entropy in Information Theory

If a random variable has distribution p , there exists an encoding with an average length of

$$H(p) \text{ bits}$$

and this is the “best” possible encoding

What happens if we use a “wrong” encoding?

- e.g. because we make an incorrect assumption on the probability distribution

If the true distribution is p , but we assume it is q , it turns out we will need to use

$$H(p) + D_{\text{KL}}(p||q) \text{ bits}$$

where $D_{\text{KL}}(p||q)$ is some measure of “distance” between p and q

Definition

The relative entropy or Kullback-Leibler (KL) divergence between two probability distributions $p(X)$ and $q(X)$ is defined as:

$$D_{\text{KL}}(p\|q) = \sum_{x \in \mathcal{X}} p(x) \log \frac{p(x)}{q(x)} = \mathbb{E}_{p(X)} \left[\log \frac{p(X)}{q(X)} \right].$$

Definition

The relative entropy or Kullback-Leibler (KL) divergence between two probability distributions $p(X)$ and $q(X)$ is defined as:

$$D_{\text{KL}}(p\|q) = \sum_{x \in \mathcal{X}} p(x) \log \frac{p(x)}{q(x)} = \mathbb{E}_{p(X)} \left[\log \frac{p(X)}{q(X)} \right].$$

- **Note:**

- ▶ Both $p(X)$ and $q(X)$ are defined over the same alphabet \mathcal{X}

- **Conventions:**

$$0 \log \frac{0}{0} \stackrel{\text{def}}{=} 0 \quad 0 \log \frac{0}{q} \stackrel{\text{def}}{=} 0 \quad p \log \frac{p}{0} \stackrel{\text{def}}{=} \infty$$

Relative Entropy

Properties

- $D_{\text{KL}}(p||q) \geq 0$

Relative Entropy

Properties

- $D_{\text{KL}}(p\|q) \geq 0$
- $D_{\text{KL}}(p\|q) = 0 \Leftrightarrow p = q$

Relative Entropy

Properties

- $D_{\text{KL}}(p\|q) \geq 0$
- $D_{\text{KL}}(p\|q) = 0 \Leftrightarrow p = q$
- $D_{\text{KL}}(p\|q) \neq D_{\text{KL}}(q\|p)$

Relative Entropy

Properties

- $D_{\text{KL}}(p\|q) \geq 0$
- $D_{\text{KL}}(p\|q) = 0 \Leftrightarrow p = q$
- $D_{\text{KL}}(p\|q) \neq D_{\text{KL}}(q\|p)$
 - ▶ Not a true distance since is not symmetric and does not satisfy the triangle inequality

Relative Entropy

Properties

- $D_{\text{KL}}(p\|q) \geq 0$
- $D_{\text{KL}}(p\|q) = 0 \Leftrightarrow p = q$
- $D_{\text{KL}}(p\|q) \neq D_{\text{KL}}(q\|p)$
 - ▶ Not a true distance since is not symmetric and does not satisfy the triangle inequality
 - ▶ Hence, “KL divergence” rather than “KL distance”

Relative Entropy

Uniform q

Let q correspond to a uniform distribution: $q(x) = \frac{1}{|\mathcal{X}|}$

Relative entropy between p and q :

$$\begin{aligned}D_{\text{KL}}(p||q) &= \sum_{x \in \mathcal{X}} p(x) \log \frac{p(x)}{q(x)} \\&= \sum_{x \in \mathcal{X}} p(x) \cdot (\log p(x) + \log |\mathcal{X}|) \\&= -H(X) + \sum_{x \in \mathcal{X}} p(x) \cdot \log |\mathcal{X}| \\&= -H(X) + \log |\mathcal{X}|.\end{aligned}$$

Matches intuition as penalty on number of bits for encoding

Relative Entropy

Example (from Cover & Thomas, 2006)

Let $X \in \{0, 1\}$ and consider the distributions $p(X)$ and $q(X)$ such that:

$$\begin{aligned} p(X = 1) &= \theta_p & p(X = 0) &= 1 - \theta_p \\ q(X = 1) &= \theta_q & q(X = 0) &= 1 - \theta_q \end{aligned}$$

What distributions are these?

Relative Entropy

Example (from Cover & Thomas, 2006)

Let $X \in \{0, 1\}$ and consider the distributions $p(X)$ and $q(X)$ such that:

$$\begin{aligned} p(X = 1) &= \theta_p & p(X = 0) &= 1 - \theta_p \\ q(X = 1) &= \theta_q & q(X = 0) &= 1 - \theta_q \end{aligned}$$

What distributions are these?

Compute $D_{\text{KL}}(p||q)$ and $D_{\text{KL}}(q||p)$ with $\theta_p = \frac{1}{2}$ and $\theta_q = \frac{1}{4}$

Relative Entropy

Example (from Cover & Thomas, 2006) — Cont'd

$$D_{\text{KL}}(p||q) = \theta_p \log \frac{\theta_p}{\theta_q} + (1 - \theta_p) \log \frac{1 - \theta_p}{1 - \theta_q}$$

Relative Entropy

Example (from Cover & Thomas, 2006) — Cont'd

$$\begin{aligned}D_{\text{KL}}(p\|q) &= \theta_p \log \frac{\theta_p}{\theta_q} + (1 - \theta_p) \log \frac{1 - \theta_p}{1 - \theta_q} \\ &= \frac{1}{2} \log \frac{\frac{1}{2}}{\frac{1}{4}} + \frac{1}{2} \log \frac{\frac{1}{2}}{\frac{3}{4}} = 1 - \frac{1}{2} \log 3 \approx 0.2075 \text{ bits}\end{aligned}$$

Relative Entropy

Example (from Cover & Thomas, 2006) — Cont'd

$$\begin{aligned}D_{\text{KL}}(p\|q) &= \theta_p \log \frac{\theta_p}{\theta_q} + (1 - \theta_p) \log \frac{1 - \theta_p}{1 - \theta_q} \\ &= \frac{1}{2} \log \frac{\frac{1}{2}}{\frac{1}{4}} + \frac{1}{2} \log \frac{\frac{1}{2}}{\frac{3}{4}} = 1 - \frac{1}{2} \log 3 \approx 0.2075 \text{ bits}\end{aligned}$$

$$D_{\text{KL}}(q\|p) = \theta_q \log \frac{\theta_q}{\theta_p} + (1 - \theta_q) \log \frac{1 - \theta_q}{1 - \theta_p}$$

Relative Entropy

Example (from Cover & Thomas, 2006) — Cont'd

$$\begin{aligned}D_{\text{KL}}(p\|q) &= \theta_p \log \frac{\theta_p}{\theta_q} + (1 - \theta_p) \log \frac{1 - \theta_p}{1 - \theta_q} \\&= \frac{1}{2} \log \frac{\frac{1}{2}}{\frac{1}{4}} + \frac{1}{2} \log \frac{\frac{1}{2}}{\frac{3}{4}} = 1 - \frac{1}{2} \log 3 \approx 0.2075 \text{ bits}\end{aligned}$$

$$\begin{aligned}D_{\text{KL}}(q\|p) &= \theta_q \log \frac{\theta_q}{\theta_p} + (1 - \theta_q) \log \frac{1 - \theta_q}{1 - \theta_p} \\&= \frac{1}{4} \log \frac{\frac{1}{4}}{\frac{1}{2}} + \frac{3}{4} \log \frac{\frac{3}{4}}{\frac{1}{2}} = -1 + \frac{3}{4} \log 3 \approx 0.1887 \text{ bits}\end{aligned}$$

- 1 Decomposability of Entropy
- 2 Relative Entropy / KL Divergence
- 3 Mutual Information**
 - Definition
 - Joint and Conditional Mutual Information
- 4 Wrapping up

Mutual Information

Definition

Let X, Y be two r.v. with joint distribution $p(X, Y)$ and marginals $p(X)$ and $p(Y)$:

Definition

The *mutual information* $I(X; Y)$ is the relative entropy between the joint distribution $p(X, Y)$ and the product distribution $p(X)p(Y)$:

$$\begin{aligned} I(X; Y) &= D_{\text{KL}}(p(X, Y) \| p(X)p(Y)) \\ &= \sum_{x \in \mathcal{X}} \sum_{y \in \mathcal{Y}} p(x, y) \log \frac{p(x, y)}{p(x)p(y)} \end{aligned}$$

Mutual Information

Definition

Let X, Y be two r.v. with joint distribution $p(X, Y)$ and marginals $p(X)$ and $p(Y)$:

Definition

The *mutual information* $I(X; Y)$ is the relative entropy between the joint distribution $p(X, Y)$ and the product distribution $p(X)p(Y)$:

$$\begin{aligned} I(X; Y) &= D_{\text{KL}}(p(X, Y) \| p(X)p(Y)) \\ &= \sum_{x \in \mathcal{X}} \sum_{y \in \mathcal{Y}} p(x, y) \log \frac{p(x, y)}{p(x)p(y)} \end{aligned}$$

Intuitively, **how much information, on average, X conveys about Y .**

Relationship between Entropy and Mutual Information

We can re-write the definition of mutual information as:

$$I(X; Y) = \sum_{x \in \mathcal{X}} \sum_{y \in \mathcal{Y}} p(x, y) \log \frac{p(x, y)}{p(x)p(y)}$$

Relationship between Entropy and Mutual Information

We can re-write the definition of mutual information as:

$$\begin{aligned} I(X; Y) &= \sum_{x \in \mathcal{X}} \sum_{y \in \mathcal{Y}} p(x, y) \log \frac{p(x, y)}{p(x)p(y)} \\ &= \sum_{x \in \mathcal{X}} \sum_{y \in \mathcal{Y}} p(x, y) \log \frac{p(x|y)}{p(x)} \end{aligned}$$

Relationship between Entropy and Mutual Information

We can re-write the definition of mutual information as:

$$\begin{aligned} I(X; Y) &= \sum_{x \in \mathcal{X}} \sum_{y \in \mathcal{Y}} p(x, y) \log \frac{p(x, y)}{p(x)p(y)} \\ &= \sum_{x \in \mathcal{X}} \sum_{y \in \mathcal{Y}} p(x, y) \log \frac{p(x|y)}{p(x)} \\ &= - \sum_{x \in \mathcal{X}} \log p(x) \sum_{y \in \mathcal{Y}} p(x, y) - \left(- \sum_{x \in \mathcal{X}} \sum_{y \in \mathcal{Y}} p(x, y) \log p(x|y) \right) \end{aligned}$$

Relationship between Entropy and Mutual Information

We can re-write the definition of mutual information as:

$$\begin{aligned} I(X; Y) &= \sum_{x \in \mathcal{X}} \sum_{y \in \mathcal{Y}} p(x, y) \log \frac{p(x, y)}{p(x)p(y)} \\ &= \sum_{x \in \mathcal{X}} \sum_{y \in \mathcal{Y}} p(x, y) \log \frac{p(x|y)}{p(x)} \\ &= - \sum_{x \in \mathcal{X}} \log p(x) \sum_{y \in \mathcal{Y}} p(x, y) - \left(- \sum_{x \in \mathcal{X}} \sum_{y \in \mathcal{Y}} p(x, y) \log p(x|y) \right) \\ &= H(X) - H(X|Y) \end{aligned}$$

Relationship between Entropy and Mutual Information

We can re-write the definition of mutual information as:

$$\begin{aligned} I(X; Y) &= \sum_{x \in \mathcal{X}} \sum_{y \in \mathcal{Y}} p(x, y) \log \frac{p(x, y)}{p(x)p(y)} \\ &= \sum_{x \in \mathcal{X}} \sum_{y \in \mathcal{Y}} p(x, y) \log \frac{p(x|y)}{p(x)} \\ &= - \sum_{x \in \mathcal{X}} \log p(x) \sum_{y \in \mathcal{Y}} p(x, y) - \left(- \sum_{x \in \mathcal{X}} \sum_{y \in \mathcal{Y}} p(x, y) \log p(x|y) \right) \\ &= H(X) - H(X|Y) \end{aligned}$$

The average reduction in uncertainty of X due to the knowledge of Y .

Mutual Information:

Properties

- Mutual Information is non-negative:

$$I(X; Y) \geq 0 \quad \text{why?}$$

Mutual Information:

Properties

- Mutual Information is non-negative:

$$I(X; Y) \geq 0 \quad \text{why?}$$

- We have seen that: $H(Y) - H(Y|X) = H(X) - H(X|Y)$, therefore:

$$I(X; Y) = I(Y; X)$$

Mutual Information:

Properties

- Mutual Information is non-negative:

$$I(X; Y) \geq 0 \quad \text{why?}$$

- We have seen that: $H(Y) - H(Y|X) = H(X) - H(X|Y)$, therefore:

$$I(X; Y) = I(Y; X)$$

- Since $H(X, Y) = H(X) + H(Y|X)$ we have that:

$$I(X; Y) = H(X) + H(Y) - H(X, Y)$$

Mutual Information:

Properties

- Mutual Information is non-negative:

$$I(X; Y) \geq 0 \quad \text{why?}$$

- We have seen that: $H(Y) - H(Y|X) = H(X) - H(X|Y)$, therefore:

$$I(X; Y) = I(Y; X)$$

- Since $H(X, Y) = H(X) + H(Y|X)$ we have that:


$$I(X; Y) = H(X) + H(Y) - H(X, Y)$$

- Finally:

$$I(X; X) = H(X) - H(X|X) = H(X)$$

Sometimes the entropy is referred to as *self-information*

Breakdown of Joint Entropy



Mutual Information

Example 1 (from Mackay, 2003)

Let X, Y, Z be r.v. with $X, Y \in \{0, 1\}$, $X \perp\!\!\!\perp Y$ and:

$$p(X = 0) = p \quad p(X = 1) = 1 - p$$

$$p(Y = 0) = q \quad p(Y = 1) = 1 - q$$

$$Z = (X + Y) \bmod 2$$

Mutual Information

Example 1 (from Mackay, 2003)

Let X, Y, Z be r.v. with $X, Y \in \{0, 1\}$, $X \perp\!\!\!\perp Y$ and:

$$p(X = 0) = p \quad p(X = 1) = 1 - p$$

$$p(Y = 0) = q \quad p(Y = 1) = 1 - q$$

$$Z = (X + Y) \bmod 2$$

(a) if $q = 1/2$ what is $P(Z = 0)$? $P(Z = 1)$? $I(Z; X)$?

Mutual Information

Example 1 (from Mackay, 2003)

Let X, Y, Z be r.v. with $X, Y \in \{0, 1\}$, $X \perp\!\!\!\perp Y$ and:

$$p(X = 0) = p \quad p(X = 1) = 1 - p$$

$$p(Y = 0) = q \quad p(Y = 1) = 1 - q$$

$$Z = (X + Y) \bmod 2$$

- (a) if $q = 1/2$ what is $P(Z = 0)$? $P(Z = 1)$? $I(Z; X)$?
- (b) For general p and q what is $P(Z = 0)$? $P(Z = 1)$? $I(Z; X)$?

Mutual Information

Example 1 (from Mackay, 2003) — Solution (a)

- (a) As $X \perp\!\!\!\perp Y$ and $q = 1/2$ the noise will flip the input with probability $q = 0.5$ regardless of the original input distribution. Therefore:

$$p(Z = 1) = \mathbb{E}[Z = 1] = 1/2 \quad p(Z = 0) = 1/2$$

Mutual Information

Example 1 (from Mackay, 2003) — Solution (a)

- (a) As $X \perp\!\!\!\perp Y$ and $q = 1/2$ the noise will flip the input with probability $q = 0.5$ regardless of the original input distribution. Therefore:

$$p(Z = 1) = \mathbb{E}[Z = 1] = 1/2 \quad p(Z = 0) = 1/2$$

Hence:

$$I(X; Z) = H(Z) - H(Z|X) = 1 - 1 = 0$$

Mutual Information

Example 1 (from Mackay, 2003) — Solution (a)

- (a) As $X \perp\!\!\!\perp Y$ and $q = 1/2$ the noise will flip the input with probability $q = 0.5$ regardless of the original input distribution. Therefore:

$$p(Z = 1) = \mathbb{E}[Z = 1] = 1/2 \quad p(Z = 0) = 1/2$$

Hence:

$$I(X; Z) = H(Z) - H(Z|X) = 1 - 1 = 0$$

Indeed for $q = 1/2$ we see that $Z \perp\!\!\!\perp X$

Mutual Information

Example 1 (from Mackay, 2003) — Solution (b)

(b)

$$\begin{aligned} \ell &\stackrel{\text{def}}{=} p(Z = 0) = p(X = 0) \times p(\text{no flip}) + p(X = 1) \times p(\text{flip}) \\ &= pq + (1 - p)(1 - q) \\ &= 1 + 2pq - q - p \end{aligned}$$

Mutual Information

Example 1 (from Mackay, 2003) — Solution (b)

(b)

$$\begin{aligned} \ell &\stackrel{\text{def}}{=} p(Z = 0) = p(X = 0) \times p(\text{no flip}) + p(X = 1) \times p(\text{flip}) \\ &= pq + (1 - p)(1 - q) \\ &= 1 + 2pq - q - p \end{aligned}$$

Similarly:

$$\begin{aligned} p(Z = 1) &= p(X = 1) \times p(\text{no flip}) + p(X = 0) \times p(\text{flip}) \\ &= (1 - p)q + p(1 - q) \\ &= q + p - 2pq \end{aligned}$$

Mutual Information

Example 1 (from Mackay, 2003) — Solution (b)

(b)

$$\begin{aligned}\ell &\stackrel{\text{def}}{=} p(Z = 0) = p(X = 0) \times p(\text{no flip}) + p(X = 1) \times p(\text{flip}) \\ &= pq + (1 - p)(1 - q) \\ &= 1 + 2pq - q - p\end{aligned}$$

Similarly:

$$\begin{aligned}p(Z = 1) &= p(X = 1) \times p(\text{no flip}) + p(X = 0) \times p(\text{flip}) \\ &= (1 - p)q + p(1 - q) \\ &= q + p - 2pq\end{aligned}$$

and:

$$\begin{aligned}I(Z; X) &= H(Z) - H(Z|X) \\ &= H(\ell, 1 - \ell) - H(q, 1 - q) \quad \text{why?}\end{aligned}$$

- 1 Decomposability of Entropy
- 2 Relative Entropy / KL Divergence
- 3 Mutual Information**
 - Definition
 - Joint and Conditional Mutual Information
- 4 Wrapping up

Joint Mutual Information

Recall that for random variables X, Y ,

$$I(X; Y) = H(X) - H(X|Y)$$

- Reduction in uncertainty in X due to knowledge of Y

More generally, for random variables $X_1, \dots, X_n, Y_1, \dots, Y_m$,

$$I(X_1, \dots, X_n; Y_1, \dots, Y_m) = H(X_1, \dots, X_n) - H(X_1, \dots, X_n | Y_1, \dots, Y_m)$$

- Reduction in uncertainty in X_1, \dots, X_n due to knowledge of Y_1, \dots, Y_m

Symmetry also generalises:

$$I(X_1, \dots, X_n; Y_1, \dots, Y_m) = I(Y_1, \dots, Y_m; X_1, \dots, X_n)$$

Conditional Mutual Information

The conditional mutual information between X and Y given $Z = z_k$:

$$I(X; Y|Z = z_k) = H(X|Z = z_k) - H(X|Y, Z = z_k).$$

Conditional Mutual Information

The conditional mutual information between X and Y given $Z = z_k$:

$$I(X; Y|Z = z_k) = H(X|Z = z_k) - H(X|Y, Z = z_k).$$

Averaging over Z we obtain:

The conditional mutual information between X and Y given Z :

$$\begin{aligned} I(X; Y|Z) &= H(X|Z) - H(X|Y, Z) \\ &= \mathbb{E}_{p(X, Y, Z)} \log \frac{p(X, Y|Z)}{p(X|Z)p(Y|Z)} \end{aligned}$$

Conditional Mutual Information

The conditional mutual information between X and Y given $Z = z_k$:

$$I(X; Y|Z = z_k) = H(X|Z = z_k) - H(X|Y, Z = z_k).$$

Averaging over Z we obtain:

The conditional mutual information between X and Y given Z :

$$\begin{aligned} I(X; Y|Z) &= H(X|Z) - H(X|Y, Z) \\ &= \mathbb{E}_{p(X, Y, Z)} \log \frac{p(X, Y|Z)}{p(X|Z)p(Y|Z)} \end{aligned}$$

The reduction in the uncertainty of X due to the knowledge of Y when Z is given.

Conditional Mutual Information

The conditional mutual information between X and Y given $Z = z_k$:

$$I(X; Y|Z = z_k) = H(X|Z = z_k) - H(X|Y, Z = z_k).$$

Averaging over Z we obtain:

The conditional mutual information between X and Y given Z :

$$\begin{aligned} I(X; Y|Z) &= H(X|Z) - H(X|Y, Z) \\ &= \mathbb{E}_{p(X, Y, Z)} \log \frac{p(X, Y|Z)}{p(X|Z)p(Y|Z)} \end{aligned}$$

The reduction in the uncertainty of X due to the knowledge of Y when Z is given.

Note that $I(X; Y; Z)$, $I(X|Y; Z)$ are illegal terms while e.g. $I(A, B; C, D|E, F)$ is legal.

- 1 Decomposability of Entropy
- 2 Relative Entropy / KL Divergence
- 3 Mutual Information
 - Definition
 - Joint and Conditional Mutual Information
- 4 Wrapping up

Summary

- Decomposability of entropy
- Relative entropy
- Mutual information
- **Reading:** Mackay §2.5, Ch 8; Cover & Thomas §2.3 to §2.5

Mutual information chain rule

Jensen's inequality

“Information cannot hurt”

Data processing inequality